
# **FRONT LABEL**



# **CONSTRUCTION**



equivalent and 0.08 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 2/days between applications. The maximum seasonal rate is 16 fl.oz. of product per 1,000 sq.ft. 1. Use Restrictions
1. Do not apply this product in a way that will contact any person or pet, either directly or Intrough drift. Keep people and pets out of the area during application.
1. Do not allow people or pets to enter the treated area until sprays have dried.
1. Apply when daytime temperatures are between 50°F and 90°F. Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur. For bermudagrass lawns do not apply when daytime temperatures exceed 85°F. 5.1 Entire Lawn Treatments Ready Spray Hose-End Sprayer

I. Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet. 2. See below for directions for the ready-to-use hose-end sprayer. 3. Spray evenly over the entire measured area Do not apply a trates above those specified on this label.
Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in Connect Shake well before using.
Connect sprayer to hose.
Turn on water. To STOP spraying, slide thumb activation switch backward to OFF position. landscape plantings and other nontarget plants.
Do not spray exposed roots of ornamentals and trees.
SHAKE WELL BEFORE USING
2. How it Works Turn off water. Relieve water pressure by sliding thumb switch to |
WATER ONLY position until water slows to a drip.
Then, slide switch back to OFF position. |• Extend hose to farthest area of the measured area to be treated. This product contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed. • To begin spraying, point nozzle in the direction you • Disconnect sprayer from hose. want to spray.

- Slide thumb activation switch forward to ON position.
• Spray evenly over measured area.
6. Cultural Tips: Warm Season Turfgrass For newly seeded areas: Bermudagrass<sup>1</sup>, common and hybrid, Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing. 6.5" For newly sodded, sprigged, or plugged areas:

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to | 4 weeks after the sodding, sprigging or plugging operations. Seeding:
- Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product. Irrigation:
Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water

6.0"

.25"

This product can be applied to the following established turfgrass species: Cool Season Turfgrass Kentucky bluegrass, Ryegrass, perennial and annual, Tall fescue, Bentgrass, Mixtures of cool season grasses containing fine fescues (red, hard and chewings)

Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed  $85^{\circ}$  F.

	<u>O NOT</u> apply this product to the following:					
	Warm Season Turfgrass	Other Areas				
St. Augustinegrass, Lawns with desir		Lawns with desirable clovers or legumes,				
	Bahiagrass,	Dichondra,				
	Centipedegrass,	Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs),				
ı	Carpetgrass,	Turfgrass species that are not listed on this label,				
	Seashore paspalum	Gardens including vegetable, fruit trees, vines and berries.				

3. Where to Use

Apply this product when weeds are small and actively growing.

**DIRECTIONS FOR USE** 

inconsistent with its labeling. I IIse Restrictions

(Crabgrass (large and smooth):

Spring: Apply this product from crabgrass emergence until the crabgrass is 3 to 4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at the same rate at 21 days later for dense populations.

| later for gense populariums.
| Foxtail (green, glant, and yellow):
| Spring and early summer: Apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at the same rate at 21 days later for dense populations. Broadleaf Weeds:

Broadlear Weeds:

Spring: Apply when dandelion, plantain, buttercup, speedwell, knotweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

Falk Apply when henbit, chickweed, white clover, ground try, lawn burweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

How Much to Use

19. How much to use
Limitations on broadcast applications to residential turfgrass
The maximum application rate is 8 fl.oz. of product per 1,000 sq.ft. per application (0.92 lb MCPA acid

| the turfgrass within 24 hours after application.


• Adequate moisture and mild environmental conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) at 24 hours before and after the application.

· Avoid applications of this product when turfgrasses are under stress since injury may result.

Avoid approximate to this product. Additional stress from Mowing:
 Delay mowing 2 days before and until 2 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
 Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

GRASS WEEDS		Crabgrass, large & smooth Foxtail, green, yellow & giant		Kikuyugrass Signalgrass, broadleaf		
BROADLEAF WEEDS						
Annual yellow sweetclover Aster Austrian fieldcress Bedstraw Beggarticks Betony, Florida Bindweed, field Bird vetch	Bittero Bitterv Black- Black I Blacks Blesse	eyed Susan	Blue lettuce Blue vervain Bracted plantain Brassbuttons Bristly oxtongue Broadleaf dock Broadleaf plantain Broomweed Buckhorn	Buckhorn plantain Bulbous buttercup Bull thistle Bullnettle Burclover Burdock Burning nettle Burwedd Buttercup		

25



3.5" -