


FRONT LABEL



<b>CONTROLS</b>	<input type="checkbox"/> NEW <input checked="" type="checkbox"/> REVISIONS <input type="checkbox"/> OTHER		approval sign-off: _____	date: _____
	client: <b>IMAGE</b>		colors: <input type="checkbox"/> CMYK <input type="checkbox"/> SPOT COLORS	
	date submitted: <b>XX/XX/17</b>	version date: <b>09/19/17</b>	requested finish: <b>XX/XX/17</b>	<input type="checkbox"/> PMS # 000 <input type="checkbox"/> PMS # 000 <input type="checkbox"/> PMS # 000 <input type="checkbox"/> PMS # 000
	file name: <b>IM68 - Image Southern Lawn Weed Killer St. Augustine &amp; Centipede Concentrate</b>		version #: <b>17</b>	notes: Special notes and instructions will go here. Special notes and instructions will go here. Special notes and instructions will go here.
	requester: _____	designer: <b>SABRINA/RYAN</b>	This color print is for content and position only. Use Contract Proof for Process-Color matching. Use Ink Drawdowns for Spot-Color matching.	

# BASE LABEL



## IMAGE<sup>®</sup>

Herbicide Consumer Concentrate

### SOUTHERN LAWN WEED KILLER

FOR ST. AUGUSTINEGRASS  
AND CENTIPEDEGRASS  
CONCENTRATE

<b>PRECAUTIONARY STATEMENTS</b>	
<b>Hazards To Humans And Domestic Animals</b>	
<b>CAUTION:</b> Causes moderate eye irritation. Harmful if absorbed through skin or if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Wash thoroughly with soap and water after handling. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.	
<b>FIRST AID</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or on clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.	

**Environmental Hazards**  
 This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to aquatic organisms in water and to plants adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.  
 This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**Manufactured for:**  
 Central Garden & Pet, Garden Division  
 1000 Parkwood Circle, Suite 700  
 Atlanta, GA 30339

1-800-265-0761  
 gardendivision@central.com  
 www.imageforweeds.com  
 EPA Reg. No. 2217-1010-90098  
 EPA Est. No. 59144-GA-1a,  
 5905-GA-01a, 5905-IA-01c,  
 5905-AR-01a  
 Subscript used is first letter of  
 run code on container.

**CENTRAL** Garden & Pet  
 Image, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.  
 ©2017 Central Garden & Pet Company

300522801-0917 • IMAGE017

.375

2.875"

Book Placement

.375

# IMAGE<sup>®</sup>

Herbicide Consumer Concentrate




## SOUTHERN LAWN WEED KILLER

FOR *ST. AUGUSTINEGRASS*  
AND *CENTPEDEGRASS*  
CONCENTRATE

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried. For any requirements specific to your state or tribe, consult the Agency responsible for pesticide regulations.


### PRODUCT FACTS:

WHAT IT DOES:	WHERE TO USE:	WHEN TO USE:
Kills major broadleaf weeds including dandelion, chickweed and clover as well as dollarweed and spurge (See inside for complete list)	Lawns: Kills the weeds not the lawn* *When used as directed	Apply when daytime temperatures are between 45° F and 90° F* *Do not apply this product to St. Augustinegrass when the temperature is above 85° F unless temporary turf injury can be tolerated.
		



**Questions & Comments:** Call 1-800-265-0761 or visit our website at [www.imageforweeds.com](http://www.imageforweeds.com).

300522691 IM68 9/17 • IM68v17



8 13576 00426 2

Manufactured for:  
Central Garden & Pet, Garden Division  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339  
[gardendivision@central.com](mailto:gardendivision@central.com)

EPA Reg. No. 2217-1010-90098  
EPA Est. No. 59144-GA-1, 5905-GA-01a,  
5905-1A-01c, 5905-AR-01b

Subscript used is first letter of run code on container.




Image, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

See inside booklet for Additional Precautionary Statements, First Aid, and complete Directions For Use.

Revise el interior para obtener declaraciones adicionales de precaución, primeros auxilios e instrucciones completas de uso.

7.85"

3.625"



**Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried. For any requirements specific to your state or tribe, consult the Agency responsible for pesticide regulations.

**1. Product Description**

Image® Herbicide Consumer Concentrate Southern Lawn Weed Killer for St. Augustinegrass and Centipedegrass Concentrate contains multiple active ingredients to provide excellent broadleaf weed control in established warm-season and cool-season turfgrasses. Weeds controlled include dandelion, chickweed, clover, dollarweed, spurge. Weed injury symptoms are normally noticed within a few days of the application and weed death can occur within 14 to 28 days. Weed injury symptoms include leaf and stem curl or twisting and weed yellowing.

**2. Use Restrictions**

- For residential use only.
- Do not apply with nozzles that produce a fine-droplet spray. Do not apply when the wind is blowing toward desirable plants. Do not spray exposed roots of ornamentals and trees.
- Do not broadcast apply to bare ground or paved surfaces.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not use on or around vegetables, fruits or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs).
- Measuring utensils such as measuring cups or measuring spoons should not be used for any food or drinking water purposes after use with this product.
- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply this product through any type of irrigation system.

**3. Where to Use**

Image® Herbicide Consumer Concentrate Southern Lawn Weed Killer for St. Augustinegrass and Centipedegrass Concentrate may be used on established lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. It will control weeds in St. Augustinegrass (including Floratam), Bermudagrass, buffalograss, centipedegrass, seashore paspalum, bahiagrass, zoysiagrass, Kentucky bluegrass, tall fescue, perennial ryegrass, and red or fine leaf fescues.

Turfgrass tolerance to this product may vary. Temporary turfgrass yellowing may occur on St. Augustinegrass and certain varieties of hybrid Bermudagrass. Do not use on bentgrass, carpetgrass, dichondra, and lawns with desirable clovers or other desirable legumes, or on ornamentals (flowers, trees, groundcovers, landscape beds, and shrubs).

**State Restrictions: New York:** Not for sale or use in New York State.

**4. How Much and When to Apply**

Mix the specified amount of product with water in a backpack, pump, hose-end, or power sprayer. Spray uniformly as a broadcast or spot treatment.

The best time for broadleaf weed applications is when weeds are young, actively growing, and soil moisture is adequate. Normally, a single application is sufficient to control broadleaf weeds in these conditions. Two applications may be necessary for hard-to-control weeds, dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for hardened-off and mature weeds.


The lower application rates within the specified range will normally provide satisfactory weed control. The higher application rates within the specified range are required for hard-to-control weeds, dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for hardened-off and mature weeds.

7.1383"

7.85"

2.875"

2.875"



**TABLE 1. Application rates for weed control in turfgrasses**

<p><b>APPLICATION SITE:</b> St. Augustinegrass (improved varieties including Floratam and Bitterblue)</p> <p><b>USE RATE PER APPLICATION:</b> Mix 5.2 to 8.6 fl. oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.</p>	<p><b>REMARKS:</b> Do not apply this product to St. Augustinegrass (all varieties) below 50° F. Do not apply to St. Augustinegrass (all varieties) when temperatures are above 85° F unless temporary injury can be tolerated. Do not make applications of this product to St. Augustinegrass when turf is going into dormancy in the fall, or if temperatures are expected to drop below 40° F within 10 days of application.</p>
<p><b>APPLICATION SITE:</b> St. Augustinegrass (common varieties)</p> <p><b>USE RATE PER APPLICATION:</b> Mix 6.2 to 8.6 fl. oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.</p>	
<p><b>APPLICATION SITE:</b> Hybrid Bermudagrass Bahigrass Zoysiagrass Buffalograss Seashore paspalum Centipedegrass</p> <p><b>USE RATE PER APPLICATION:</b> Mix 6.2 to 8.2 fl. oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.</p>	<p><b>REMARKS:</b> This product may be applied to dormant bermudagrass, dormant zoysiagrass, and dormant bahigrass. Use caution when applying this product to warm-season turfgrasses when turf is going into dormancy in the fall or coming out of dormancy in the spring as some injury may occur.</p>

<p><b>APPLICATION SITE:</b> Common Bermudagrass</p> <p><b>USE RATE PER APPLICATION:</b> Mix 7.2 to 8.6 fl. oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.</p>	<p><b>REMARKS:</b> This product may be applied to dormant bermudagrass, dormant zoysiagrass, and dormant bahigrass. Use caution when applying this product to warm-season turfgrasses when turf is going into dormancy in the fall or coming out of dormancy in the spring as some injury may occur.</p>
<p><b>APPLICATION SITE:</b> Kentucky Bluegrass Annual bluegrass</p> <p><b>USE RATE PER APPLICATION:</b> Mix 7.2 to 10.6 fl. oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.</p>	<p><b>REMARKS:</b> In mixed stands use 4.8 fl. oz. per 1000 sq. ft. unless some injury to the sensitive species can be tolerated.</p>
<p><b>APPLICATION SITE:</b> Perennial ryegrass Annual ryegrass Tall fescue Red or fine leaf fescues</p> <p><b>USE RATE PER APPLICATION:</b> Mix 4.8 fl. oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.</p>	

**Limitations on Treatments:**  
Maximum rate per application is 10.6 fl.oz. per 1000 sq.ft. [0.40 lb 2,4-D ae per acre, 0.046 lb penoxsulam per acre, 0.11 lb dicamba ae per acre, and 0.046 lb sulfentrazone per acre]. Maximum rate per season is 21.2 fl.oz. per 1000 sq.ft. [0.80 lb 2,4-D ae per acre, 0.092 lb penoxsulam per acre, 0.22 lb dicamba ae per acre, and 0.092 lb sulfentrazone per acre]. Do not apply more than 2 broadcast treatments per season. A minimum of 30 days is required between applications.

2.375"

2.375"

### 5. How to Apply Hand-Operated Tank Sprayers: For hand-operated sprayers (backpack, knapsack, tank-type or pump-up):

- Measure the total lawn area to be sprayed.  
For rectangular areas, multiply the length by the width to obtain total square feet.
- Mix 2 2/3 fl. oz. per gallon of water for every 500 sq.ft.
- Spray evenly over the lawn. Spray coverage should be uniform and complete.

### 6. Spray Drift Management

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash from treated areas onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

### 7. Use Precautions

- Temporary yellowing may occur to some turfgrasses following application. Sensitive species include St. Augustinegrass, Centipedegrass and hybrid Bermudagrass.
- Applications made when ambient temperatures are above 90°F (85°F for St. Augustinegrass) may cause undesirable turf injury. Do not apply at these temperatures unless temporary turf injury can be tolerated.
- Rainfast in as little as 4 hours. Do not apply this product immediately before rainfall or irrigation. Do not irrigate or water the turfgrass within 3 hours after application. If dry conditions exist, irrigate at least 8 hours before and 8 hours after application.
- Delay mowing 2 days before and until 2 days after the application of this product.
- To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Injury can occur if this product is applied under any of these or other stress conditions.
- Do not apply this product to warm-season turfgrasses when turf is going into dormancy in the fall or coming out of dormancy in the spring as some injury may occur.
- Avoid fall or winter applications unless turf stunting can be tolerated.

### 8. Broadleaf Weeds Controlled

This product will control or suppress the following broadleaf. Apply when weeds are young and actively growing.

#### BROADLEAF WEEDS


Aster, white heath & white prairie	Dollarweed (pennywort)
Bedstraw	Doveweed*
Beggarweed, creeping	English daisy
Beggarweed, dwarf	False dandelion (spotted catsear & common catsear)
Bindweed	Field bindweed (morningglory & creeping jenny)
Black medic	Field oxeye-daisy (creeping oxeye)
Broadleaf plantain	Filaree, whitestem & redstem
Bull thistle	Florida betony (rattlesnake weed)*
Burclover	Florida pusley
Burdock, common	Goldenrod
Buttercup, creeping	Ground ivy
Carpetweed	Groundsel
Catsear, spotted	Hawkweed
Chamberbitter*	Healall
Chickweed, common	Henbit
Chickweed, mouseear	Innocence (Blue-eyed Mary)
Chicory	Knotweed
Cinquefoil	Lambsquarters
Clover	Lawn burweed
Clover, hop	Lespedeza, common
Clover, red	Mallow, common
Clover, white (Dutch clover, honeysuckle clover, white trefoil, & purplewort)	Matchweed
Cocklebur	Nutsedge (yellow)
Coffeeweed	Old world diamond flower
Compassplant	
Curly dock	
Dandelion	
Dayflower	
Deadnettle	
Dock	
Dogfennel	

7.1383"

7.85"

2.875"

2.875"



**BROADLEAF WEEDS**

Oxalis (yellow woodsorrel & creeping woodsorrel)	Smartweed
Parsley-piert	Spurge
Pennsylvania smartweed	Spurweed
Pepperweed	Thistle
Pigweed	Veronica (corn speedwell)*
Pineappleweed	Vetch, common
Plantain, buckhorn*	Virginia buttonweed *
Plantain, common*	Virginia pepperweed
Poison ivy	Wild carrot
Poison oak	Wild garlic
Puncturevine	Wild geranium
Purple cudweed	Wild lettuce
Purslane	Wild mustard
Ragweed	Wild onion
Redweed	Wild strawberry *
Red sorrel (sheep sorrel)	Wild violet *
Roundleaf mallow	Yarrow
Shepherd's purse	Yellow rocket

\* Hard-to-control: May require two applications and higher use rates within the specified rate range.

**STORAGE AND DISPOSAL**  
Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in the original container in a locked storage area. Keep from freezing.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**  
FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

**NOTICE:** To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

If for any reason you are not satisfied after using this product, simply send us original proof of purchase and we will refund the purchase price.

**Questions or comments?**  
Call toll free 1-800-265-0761.

