

WON'T HARM
LAWNS*

*when used as directed

IMAGE[®]
HERBICIDES

WEED KILLER

FOR SOUTHERN LAWNS

CONCENTRATE

For use on Bermuda, St. Augustinegrass,
Zoysia, and Centipedegrass



Dandelion



Chickweed



Clover



Dollarweed

**KILLS MAJOR BROADLEAF WEEDS,
ROOTS AND ALL**

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See attached booklet for
complete Precautionary
Statements and Directions for Use, including Storage
and Disposal, and back panel for First Aid.

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL
PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

300535848-05/24 | IM216 V9

ACTIVE INGREDIENTS:

Penoxsulam	0.14%
Sulfentrazone	0.14%
2,4-D, dimethylamine salt	1.46%
Dicamba, dimethylamine salt	0.41%

OTHER INGREDIENTS: 92.85%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

0.013 lb Penoxsulam per gallon or 0.14%
0.013 lb Sulfentrazone per gallon or 0.14%
0.109 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 1.21%
0.031 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.34%

32 FL OZ COVERS UP TO

6,670 SQ. FT.

NET CONTENTS

1 QUART

(32 FL OZ) (946 mL)



WEED KILLER

FOR SOUTHERN LAWNS

CONCENTRATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin or if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Wash thoroughly with soap and water after handling. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

FIRST AID

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to aquatic organisms in water and to plants adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

GROUNDWATER ADVISORY STATEMENT: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

SURFACE WATER ADVISORY STATEMENT: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of penoxsulam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 2217-1010-90098
EPA Est. No. 59144-GA-1, 92162-MD-1,
Subscript used is first letter of run code on container.

1-800-265-0761
gardendivision@central.com
www.imageforweeds.com

Image, Central Garden & Pet and Central Garden & Pet with
design are trademarks of Central Garden & Pet Company.

© 2017-2024 Central Garden & Pet Company

300535849 - 5/24 | IM216 V9

[illegible]

Kills major broadleaf weeds including dandelion, chickweed and clover as well as dollarweed and spurge
(See inside for complete list)



Lawns: Kills the weeds not the lawn*
* When used as directed



Apply when daytime temperatures are between 50° F and 90° F*
 * Do not apply this product to St. Augustinegrass when the temperature is above 85° F unless temporary turf injury can be tolerated.



Call 1-800-265-0761 or visit our website at www.imageforweeds.com

See attached booklet for complete Precautionary Statements and Directions for Use, including Storage and Disposal, and back panel for First Aid.



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 2217-1010-90098
EPA Est. No. 59144-GA-I, 92162-MO-I
Subscript used is first letter of run code on container.
1-800-265-0761
gardendivision@central.com

300535849-5/24 | IM216 V9



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation, Harmful if absorbed through skin or if swallowed, When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves, Wash thoroughly with soap and water after handling, After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water, Remove saturated clothing as soon as possible and shower,

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment, You may also contact 1-800-265-0761 for emergency medical treatment information.	

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark, Drift or runoff may be hazardous to aquatic organisms in water and to plants adjacent to treated areas, Do not contaminate water when disposing of equipment wash waters or rinsate, To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters, Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area, Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems,

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site, Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift,

GROUNDWATER ADVISORY STATEMENT: This chemical has properties and characteristics associated with chemicals detected in groundwater, This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow,

SURFACE WATER ADVISORY STATEMENT: This product may impact surface water quality due to runoff of rain water, This is especially true for poorly draining soils and soils with shallow ground water, This product is classified as having high potential for reaching surface water via runoff for several months or more after application, A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of penoxsulam from runoff water and sediment, Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours,

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling, Do not apply this product in a way that will contact any person or pet, either directly or through drift, Keep people and pets out of the area during application, Do not allow people or pets to enter the treated area until sprays have dried, For any requirements specific to your state or tribe, consult the Agency responsible for pesticide regulations,

1. Product Description

This product contains multiple active ingredients to provide excellent broadleaf weed control in established warm-season and cool-season turfgrasses, Weeds controlled include dandelion, chickweed, clover, dollarweed, spurge, Weed injury symptoms are normally noticed within a few days of the application and weed death can occur within 14 to 28 days, Weed injury symptoms include leaf and stem curl or twisting and weed yellowing,

2. Use Restrictions

- For residential use only,
- Do not apply with nozzles that produce a fine-droplet spray.
- Do not apply when the wind is blowing toward desirable plants.
- Do not spray exposed roots of ornamentals and trees,
- Do not broadcast apply to bare ground or paved surfaces,
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens,
- Do not use on or around vegetables, fruits or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs),
- Measuring utensils such as measuring cups or measuring spoons should not be used for any food or drinking water purposes after use with this product,
- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays), Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays),
- Do not apply this product through any type of irrigation system,

3. Where to Use

This product may be used on established lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present, It will control weeds in St. Augustinegrass (including Floratam), Bermudagrass, buffalograss, centipedegrass, seashore paspalum, bahiagrass, zoysiagrass, Kentucky bluegrass, tall fescue,

perennial ryegrass, and red or fineleaf fescues, Turfgrass tolerance to this product may vary, Temporary turfgrass yellowing may occur on St. Augustinegrass and certain varieties of hybrid Bermudagrass, Do not use on bentgrass, carpetgrass, dichondra, and lawns with desirable clovers or other desirable legumes, or on ornamentals (flowers, trees, groundcovers, landscape beds, and shrubs),

State Restrictions: New York: Not for sale or use in New York State.

4. How Much and When to Apply

Mix the specified amount of product with water in a backpack, pump, hose-end, or power sprayer, Spray uniformly as a broadcast or spot treatment, The best time for broadleaf weed applications is when weeds are young, actively growing, and soil moisture is adequate, Normally a single application is sufficient to control broadleaf weeds in these conditions, Two applications may be necessary for hard-to-control weeds, dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for hardened-off and mature weeds, The lower application rates within the specified range will normally provide satisfactory weed control, The higher application rates within the specified range are required for hard-to-control weeds, dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for hardened-off and mature weeds,

TABLE 1. Application rates for weed control in turfgrasses	
APPLICATION SITE: St. Augustinegrass (improved varieties including Floratam and Bitterblue)	REMARKS: Do not apply this product to St. Augustinegrass (all varieties) below 50° F. Do not apply to St. Augustinegrass (all varieties) when temperatures are above 85° F unless temporary injury can be tolerated. Do not make applications of this product to St. Augustinegrass when turf is going into dormancy in the fall, or if temperatures are expected to drop below 40° F within 10 days of application.
USE RATE PER APPLICATION: Mix 5.2 to 8.6 fl.oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.	
APPLICATION SITE: St. Augustinegrass (common varieties)	
USE RATE PER APPLICATION: Mix 6.2 to 8.6 fl.oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.	
APPLICATION SITE: Hybrid Bermudagrass Bahiagrass Zoysiagrass Buffalograss Seashore paspalum Centipedegrass	REMARKS: This product may be applied to dormant bermudagrass, dormant zoysiagrass, and dormant bahiagrass. Use caution when applying this product to warm-season turfgrasses when turf is going into dormancy in the fall or coming out of dormancy in the spring as some injury may occur.
USE RATE PER APPLICATION: Mix 6.2 to 8.2 fl.oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.	
APPLICATION SITE: Common Bermudagrass	
USE RATE PER APPLICATION: Mix 7.2 to 8.6 fl.oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.	
APPLICATION SITE: Kentucky bluegrass Annual bluegrass	REMARKS: In mixed stands use 4.8 fl.oz. per 1000 sq. ft. unless some injury to the sensitive species can be tolerated.
USE RATE PER APPLICATION: Mix 7.2 to 10.6 fl.oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.	
APPLICATION SITE: Perennial ryegrass Annual ryegrass Tall fescue Red or fine leaf fescues	
USE RATE PER APPLICATION: Mix 4.8 fl.oz. in 1 to 4 gallons of water and spray over 1000 sq. ft.	

Limitations on Treatments:

Maximum rate per application is 10.6 fl.oz. per 1000 sq. ft. (0.40 lb 2,4-D ae per acre, 0.046 lb penoxsulam per acre, 0.31 lb dicamba ae per acre, and 0.046 lb sulfentrazone per acre), Maximum rate per season is 21.2 fl.oz. per 1000 sq. ft. (0.80 lb 2,4-D ae per acre, 0.092 lb penoxsulam per acre, 0.22 lb dicamba ae per acre, and 0.092 lb sulfentrazone per acre), Do not apply more than 2 broadcast treatments per season, A minimum of 30 days is required between applications,

5. How to Apply

Hand-Operated Tank Sprayers

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up):

- Measure the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- Measure the specified amount of product and mix with water in the sprayer.
- Spray evenly over the lawn. Spray coverage should be uniform and complete.

6. Spray Drift Management

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash from treated areas onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

7. Use Precautions

- Temporary yellowing may occur to some turfgrasses following application. Sensitive species include St. Augustinegrass, Centipedegrass and hybrid Bermudagrass.
- Applications made when ambient temperatures are above 90°F (85°F for St. Augustinegrass) may cause undesirable turf injury. Do not apply at these temperatures unless temporary turf injury can be tolerated.
- Rainfast in as little as 4 hours. Do not apply this product immediately before rainfall or irrigation. Do not irrigate or water the turfgrass within 3 hours after application. If dry conditions exist, irrigate at least 8 hours before and 8 hours after application.
- Delay mowing 2 days before and until 2 days after the application of this product.
- To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Injury can occur if this product is applied under any of these or other stress conditions.
- Do not apply this product to warm-season turfgrasses when turf is going into dormancy in the fall or coming out of dormancy in the spring as some injury may occur.
- Avoid fall or winter applications unless turf stunting can be tolerated.

8. Broadleaf Weeds Controlled

This product will control or suppress the following broadleaf weeds. Apply when weeds are young and actively growing.

BROADLEAF WEEDS		
Aster, white heath & white prairie	Doveweed*	Pigweed
Bedstraw	English daisy	Pineappleweed
Beggarweed, creeping	False dandelion (spotted	Plantain, buckhorn*
Beggarweed, dwarf	catsear & common	Plantain, common*
Bindweed	catsear)	Poison ivy
Black medic	Field bindweed	Poison oak
Broadleaf plantain	(morningglory & creeping	Puncturevine
Bull thistle	jenny)	Purple cudweed
Burdock	Field oxeye-daisy (creeping	Purslane
Burdock, common	oxeye)	Ragweed
Buttercup, creeping	Flaree, whitestem & redstem	Redweed
Carpetweed	Florida betony (rattlesnake	Red sorrel (sheep sorrel)
Catsear, spotted	weed)*	Roundleaf mallow
Chamberbitter*	Florida pusley	Shepherd's purse
Chickweed, common	Goldenrod	Smartweed
Chickweed, mouseear	Ground ivy	Spurge
Chicory	Groundsel	Spurweed
Cinquefoil	Hawkweed	Thistle
Clover	Heckall	Veronica (corn
Clover, hop	Henbit	speedwell)*
Clover, red	Innocence (blue-eyed Mary)	Vetch, common
Clover, white (Dutch clover,	Knotweed	Virginia buttonweed *
honeyuckle clover, white	Lambsquarters	Virginia pepperweed
trefoil, & purplewort)	Lawn burweed	Wild carrot
Cockspur	Lespedeza, common	Wild garlic
Coffeeweed	Mallow, common	Wild geranium
Compassplant	Matchweed	Wild lettuce
Cutly dock	Nutsedge (yellow)	Wild mustard
Dandelion	Old world diamond flower	Wild onion
Dayflower	Oxalis (yellow woodsorrel &	Wild strawberry *
Deadnettle	creeping woodsorrel)	Wild violet *
Dock	Parsley-piert	Yarrow
Dogfennel	Pennsylvania smartweed	Yellow rocket
Dollarweed (pennywort)	Pepperweed	
* Hard-to-control: May require two applications and higher use rates within the specified rate range.		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions. If for any reason you are not satisfied after using this product, simply send us original proof of purchase and we will refund the purchase price.

Questions or comments?

Call toll free 1-800-265-0761.

Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

NOTES

[illegible]

NOTES

[illegible]