

COLOR KEY

- **Dieline**
- **LABELS: Safe Copy Area**
All copy and printable items must be within this area so they are not trimmed off during production. Items **MUST** stay inset -0.0625 (1/16") away from the dieline edge, excluding graphics intended to bleed.
- **BOOKS: Safe Copy Area**
- **Front:** 125" inset from edge of die
- **All inside pages:** .25" inset from edge of die
- **Base:** .25" inset from the book placement
- **Required Bleed** - We **REQUIRE** at least 0.25" (1/4") of bleed past the dieline for any background, including images.

PLEASE NOTE:

- 1.) SPECIFY ALL COLORS TO BE USED as spot Pantone colors **or** 4 color process (CMYK).
- 2.) Please drag ALL SPOT COLORS utilized into the SWATCHES pallet.
- 3.) All black items should use the SAME BLACK SWATCH in the color pallet, 100% Black.

FILES:

Adobe Illustrator (.ai) files are best suited for label production and design.

DIELINE TEMPLATE

Date: 2/3/22
Size: 6.6 x 4.39
Artist: CD
RLG#: 12127-609429

CONTROLS	<input checked="" type="checkbox"/> NEW <input type="checkbox"/> REVISIONS <input type="checkbox"/> OTHER			approval sign-off: _____ date: _____	
	client: IMAGE			<input checked="" type="checkbox"/> CMYK <input checked="" type="checkbox"/> SPOT COLORS	
	date submitted: XX/XX/21	version date: 07/26/24	requested finish: XX/XX/21	<input type="checkbox"/> PMS # 715	<input type="checkbox"/> PMS # 7546
	file name: IM213 - IMAGE 300535842 Crabgrass_Weed Killer 320Z RTS			<input type="checkbox"/> PMS # 9062	<input type="checkbox"/> PMS # 000
	requester: KATHLEEN	designer: MICHELLE/BRENT	version #: 8		

This color print is for content and position only. Use Contract Proof for Process-Color matching. Use Ink Drawdowns for Spot-Color matching.

CENTRAL
Garden & Pet
CENTRAL GARDEN & PET | v2.0 | 42/13

COLOR KEY

- **Dieline**
- **LABELS: Safe Copy Area**
All copy and printable items must be within this area so they are not trimmed off during production. Items **MUST** stay inset -0.0625 (1/16") away from the dieline edge, excluding graphics intended to bleed.
- **BOOKS: Safe Copy Area**
 - **Front:** 125" inset from edge of die
 - **All inside pages:** .25" inset from edge of die
 - **Base:** .25" inset from the book placement
- **Required Bleed** - We REQUIRE at least 0.25" (1/4") of bleed past the dieline for any background, including images.

PLEASE NOTE:

- 1.) SPECIFY ALL COLORS TO BE USED as spot Pantone colors **or** 4 color process (CMYK).
- 2.) Please drag ALL SPOT COLORS utilized into the SWATCHES pallet.
- 3.) All black items should use the SAME BLACK SWATCH in the color pallet, 100% Black.

FILES:

Adobe Illustrator (.ai) files are best suited for label production and design.

DIELINE TEMPLATE

Date: 2/3/22
Size: 6.48 X 3.13
Artist: CD
RLG#: 12127-609430

IMAGE
HERBICIDES
**CRABGRASS
& WEED KILLER**
FOR LAWN
READY-TO-SPRAY

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment advice.</p>	

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to waters. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment wash waters or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 2217-954-90098
EPA Est. No. 59144-GA-1, 92162-MO-1
Subscript used is first letter of run code on container.

1-800-265-0761
gardendivision@central.com
www.imageforweeds.com
Image, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.
©2017-2024 Central Garden & Pet Company

300535843 - 7/24 - IM213 v8

STARTS WORKING
IMMEDIATELY

IMAGE[®]
HERBICIDES

CRABGRASS & WEED KILLER

FOR LAWN

READY-TO-SPRAY

PRODUCT FACTS:



WHAT IT DOES:

Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See inside for complete list)



WHERE TO USE:

Lawns: Kills the weeds not the lawn when used as directed. This product can be applied to the following established turfgrass species:

Cool season turfgrass: Kentucky bluegrass, Ryegrass, perennial and annual, tall fescue, Bentgrass, Mixtures of cool season grasses containing fine fescues (red, hard, and chewings)

Warm season turfgrass: Bermudagrass¹, common and hybrid, Zoysiagrass, Buffalograss

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.



WHEN TO USE:

Apply when daytime temperatures are between 50° F and 90° F. (Except on Bermudagrass - Do not apply when daytime temperatures exceed 85° F)



Questions & Comments:

Call 1-800-265-0761 or visit our website at www.imageforweeds.com

See attached booklet for complete Precautionary Statements and Directions for Use, including Storage and Disposal, and back panel for First Aid



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 2217-954-90098
EPA Est. No. 59144-BA-1, 52162-MD-1,
Subscript used is first letter of run code
on container.

1-800-265-0761
gardendivision@central.com

300535843-7/24-IM213 v8



8 13576 00410 1

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Use Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Apply when daytime temperatures are between 50° F and 90° F. Do not treat when air temperatures exceed 90° F as damage to turfgrass may occur. For bermudagrass lawns do not apply when daytime temperatures exceed 85° F.
- Do not apply at rates above those specified on this label.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

SHAKE WELL BEFORE USING

2. How it Works

This product contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

3. Where to Use

This product can be applied to the following established turfgrass species:	
Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass, Ryegrass, perennial and annual, Tall fescue, Bentgrass, Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	Bermudagrass, common and hybrid, Zoysiagrass, Buffalograss
Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.	
DO NOT apply this product to the following:	
Warm Season Turfgrass	Other Areas
St. Augustinegrass, Bahiagrass, Centipede grass, Carpetgrass, Seashore paspalum	Lawns with desirable clovers or legumes, Dichondra, Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Turfgrass species that are not listed on this label, Gardens including vegetables, fruit trees, vines and berries.

4. When to Use:

Apply this product when weeds are small and actively growing. Crabgrass (large and smooth):

- **Spring:** Apply this product from crabgrass emergence until the crabgrass is 3 to 4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at the same rate at 21 days later for dense populations.

Foxtail (green, giant, and yellow):

- **Spring and early summer:** Apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at the same rate at 21 days later for dense populations.

Broadleaf Weeds:

- **Spring:** Apply when dandelion, plantain, buttercup, speedwell, knotweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- **Fall:** Apply when henbit, chickweed, white clover, ground ivy, lawn burweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

5. How Much to Use

Limitations on broadcast applications to residential turfgrass

The maximum application rate is 6 fl.oz. of product per 1,000 sq.ft. per application (0.92 lb MCPA acid equivalent and 0.08 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 16 fl.oz. of product per 1,000 sq.ft. (1.84 lb MCPA ae and 0.17 lb dicamba ae per acre acid equivalent per acre).

5.1 Entire Lawn Treatments

Ready Spray Hose-End Sprayer

1. Measure the entire area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
2. See below for directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the entire measured area.

CONNECT

1. Shake well before use.
2. Connect sprayer to garden hose.
3. Turn on water.

SPRAY

1. Extend hose to farthest area to be treated, point nozzle in direction to be sprayed and turn on sprayer. Work back toward the faucet to avoid walking over treated grass.
2. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas.

FINISH

1. Turn off sprayer.
2. Turn off water at faucet.
3. Relieve pressure in hose by turning water back on at sprayer until water slows to a drip. Close sprayer. Disconnect sprayer from hose.

6. Cultural Tips:

For newly seeded areas:

- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged, or plugged areas:

- The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging or plugging operations.

Seeding:

- Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- Adequate moisture and mild environmental conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) at 24 hours before and after the application.
- Avoid applications of this product when turfgrasses are under stress since injury may result.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

7. Weeds Controlled

GRASS WEEDS	Crabgrass, large & smooth Foxtail, green, yellow & giant	Kikuyugrass Signalgrass, broadleaf
BROADLEAF WEEDS		
Annual yellow sweetclover	Elderberry	Red sorrel
Aster	False dandelion	Redroot pigweed
Austrian fieldcress	False flax	Redstem filaree
Bedstraw	False sunflower	Rough cinquefoil
Beggarticks	Fiddleneck	Rough fleabane
Betony, Florida	Florida pusley	Russian pigweed
Bindweed, field	Frenchweed	Russian thistle
Bird vetch	Gallinsoga	Scarlet pimpernel
Bitter wintercress	Goathead	Scotch thistle
Bittercress, hairy	Goldenrod	Sheep sorrel
Bitterweed	Ground Ivy	Shepherdspurse
Black-eyed Susan	Gumweed	Slender plantain
Black medic	Hairy fleabane	Smallflower gallinsoga
Black mustard	Hawkweed	Smooth dock
Blackseed plantain	Healall	Smooth pigweed
Blessed thistle	Heartleaf drymary	Sorrel
Bloodflower	Heathaster	Sowthistle
milkweed	Hedge bindweed	Spanishneedles
Blue lettuce	Hedge mustard	Speedwell
Blue vervain	Hemp	Spiny amaranth
Bracted plantain	Henbit	Spiny cocklebur
Brassbuttons	Hoary cress	Spiny sowthistle
Bristly oxtongue	Hoary plantain	Spotted catsear
Broadleaf dock	Hoary vervain	Spurred spurge
Broadleaf plantain	Horsenettle	Spurweed
Broomweed	Jimsonweed	Stinging nettle
Buckhorn	Knawel	Strawberry, India mock
Buckhorn plantain	Knotweed	Tall nettle
Bulbous buttercup	Kochia	Tall vervain
Bull thistle	Lambsquarters	Tansy ragwort
Bullnettle	Lespedeza	Tansy mustard
Burdock	Mallow	Tanweed
Burning nettle	Matchweed	Thistle
Burweed	Mexicanweed	Trailing crownvetch
Buttercup	Milk vetch	Tumble mustard
Buttonweed	Momingglory	Tumble pigweed
Canada thistle	Mouseear hawkweed	Velvetleaf
Carolina geranium	Mugwort	Venice mallow
Carpetweed	Musk thistle	Virginia buttonweed
Catchweed bedstraw	Mustard	Virginia creeper
Catnip	Narrowleaf plantain	Virginia pepperweed
Catsear	Narrowleaf vetch	Wavyleaf bullthistle
Chickweed, common	Nettle	Western daisy
Chickweed, mouseear	Orange hawkweed	Western salisfly
Chicory	Oriental cocklebur	White mustard
Cinquefoil	Oxalis	Wild aster
Clover, crimson	Parsley-piert	Wild buckwheat
Clover, hop	Parsnip	Wild carrot
Clover, red	Pearlwort	Wild four-o'clock
Clover, strawberry	Pennywort	Wild garlic
Clover, sweet	Pennywort	Wild geranium
Clover, white	Peppergrass	Wild lettuce
Cockle	Pepperweed	Wild marigold
Cocklebur	Pigweed	Wild mustard
Common mullein	Pineywoods bedstraw	Wild onion
Creeping Jenny (Creeping Charlie)	Plains coreopsis (tickseed)	Wild parsnip
Cudweed	Plantain	Wild radish
Curly dock	Poison ivy	Wild rape
Daisy, English	Poison oak	Wild strawberry
Daisy, fleabane	Pokeweed	Wild sweet potato
Daisy, oxeye	Poorjoe	Wild vetch
Dandelion	Prairie sunflower	Wild violet
Dichondra	Prickly lettuce	Woodsorrel
Dogbane	Prickly sida	Woody croton
Dogfennel	Prostrate knotweed	Woody plantain
Dollarweed	Prostrate pigweed	Wormseed
	Prostrate spurge	Yarrow
	Prostrate vervain	Yellow rocket
	Puncturevine	Yellowflower pepperweed
	Purslane, common	
	Ragweed	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

